ACP American College of Physicians Leading Internal Medicine, Improving Lives | Minnesota Chapter

Medical Students

Research - Medical Students

Kristene Tadese
Dr. LaPrincess Brewer
Sarah Jenkins

Factors Facilitating Academic-Community Research Partnerships with African-American Churches

Background: African-Americans (AAs) continue to be underrepresented in medical research and clinical trials. This can lead to inadequate access to novel medical therapies and innovations among this population, thus undermining health equity. AA churches are valuable partners in implementing health promotion interventions to combat disparities. This study evaluates church characteristics associated with enrollment into the FAITH! Trial, a community-based, cluster randomized trial for cardiovascular health (CVH) promotion among AA churches.
Methods: Churches attending recruitment events in Minneapolis-St. Paul and Rochester, MN were invited to complete an electronic survey and telephone interview. These instruments assessed church characteristics and readiness to engage in health promotion programming (HPP) as determined by the Brand's PREACH (Predicting Readiness to Engage African American Churches in Health) assessment tool. The primary outcome of interest was church enrollment in the FAITH! Trial. Key predictors of interest included congregation size, average congregation member age and church readiness to engage in HPP as defined by PREACH scores and stage classifications (1-3), denoting low to high capacity for engagement.

Results: Of the 30 churches attending recruitment events, 26 (93%) completed the screening survey and follow-up interview, and 16 (53%) enrolled into the

	intervention study had greater infrastructure for HPP, larger congregations and members of younger age. Institutions seeking to increase accessibility to clinical trials and medical therapies to AA communities by partnering with AA churches must provide greater support to smaller congregations with members of older age through adequate personnel and funding for optimal outcomes.
Christina Gu Dr. Patricia Jewett Dr. K. Robin Yabroff Dr. Rachel Vogel Dr. Helen Parsons Dr. Himal Purani Dr. Anne Blaes	*Regional Clustering of Financial Hardship in the United States Among Cancer Survivors by Age, Sex, and Race/Ethnicity* Background: Innovative treatments have improved cancer survival but also increased financial hardship for patients. While demographic factors associated with financial hardship among cancer survivors are known in the United States (US), the role of geography is less clear. Methods: We evaluated financial hardship (self-reported foregoing care due to cost within 12 months) by US Census region (Northeast, North Central/Midwest [NCMW], South, and West) in combination with demographic factors (age, sex, race/ethnicity) among 217,981 adult cancer survivors aged 20 to 82 years who participated in the 2015-2019 Behavioral Risk Factor Surveillance System survey. We summarized region-and group-specific prevalence of medical financial hardship and used multivariable logistic regression models to compare financial hardship by region. Results: Financial hardship was most prevalent in the South (19-38% among those aged <65 years; 4-

autoimmune processes (C3, C4, CH50, anti-

Case Presentation: A seventy-

the hospital, and can lead to life threatening sepsis. Cuts, scrapes, and puncture wounds are all potential entry points for an infection and can be caused by a variety of traumatic mechanisms that disrupt the skin barrier. Diabetics with neuropathy and poor vascularity are at increased risk of unidentified foot wounds, but the hands also remain at risk for injury secondary to activities of daily living.

Case Presentation: A 69 year old male with history of uncontrolled type 2 diabetes mellitus, hypertension, atrial fibrillation, and bilateral transtibial amputations secondary to chronic foot ulcers, presented to the hospital with altered mental status. Non-con CT of the head did not reveal acute intracranial hemorrhage, electrolytes within normal limits, but labs were notable for elevated white count, CRP, ESR, lactate, and hyperglycemia. Patient was febrile, tachycardic with irregularly irregular rhythm. Right thumb markedly swollen, erythematous, and warm with redness extending into the thenar eminence and paronychia. The patient is unable to recall any injury to his thumb. A clinical diagnosis of sepsis secondary to right thumb cellulitis was made. Blood cultures were obtained and broad-spectrum antibiotic coverage with vancomycin and zosyn was started. Xray of the thumb was negative for gas, foreign body, or osteomyelitis. Orthopedic surgeons performed bedside

Methods: The patient received her first infusion of pembrolizumab, an immune checkpoint inhibitor (ICI), two weeks prior to presentation and) after progression of her malignancy on other therapies. She presented to the emergency depar

106 beats per minute, respiratory rate 19 to 30 per minute, and saturating 89%

	Methods: Echocardiogram revealed moderately reduced global systolic function with an estimated EF of 35% and multiple segmental wall motion abnormalities, suggestive of takotsubo cardiomyopathy. Coronary CT angiography showed patent coronary arteries with no evidence of an ischemic event. She was medically managed for her takotsubo cardiomyopathy with beta blockers, and isosorbide dinitrite in addition to her chronic aspirin, ranolazine, and rosuvastatin for comorbid CAD. She improved and was discharged to home after three days. One month post discharge her echocardiogram was normal with no wall motion abnormalities and EF of 55%. documented episode of takotsubo cardiomyopathy. She reported that she typically has one to two episodes of cardiomyopathy per year that are linked is the highest number of recurrent episodes of takotsubo cardiomyopathy documented in the medical literature. Conclusion: Takotsubo cardiomyopathy is an important clinical diagnosis whose incidence is increasing. A JAMA study found that incidence of takotsubo cardiomyopathy has increased by roughly 8% in the setting of the COVID-19 pandemic. Takotsubo cardiomyopathy disproportionately impacts women and one study found 55.8% of patients with takotsubo cardiomyopathy had psychiatric or neurological disorders compared to 25% of patients with ACS. This case demonstrates that although stress cardiomyopathy recurrence is rare, it should be included in the differential. Additionally, these patients should have multifactorial management that treats not only the physiological symptoms of the cardiomyopathy, but also targets the psychological stressors that caused the pathology. Proper management of
Anja Touma Dr. Georgios Manousakis Dr. Prabhjot Nijjar	*A Rare Case of Severe Cardiomyopathy Associated with Myotonic Dystrophy Type 2* Background: Myotonic dystrophies (DM) are multisystemic diseases characterized by muscle weakness and myotonia. DM2 (myotonic dystrophy Type 2; proximal myotonic myopathy; PROMM) is an autosomal dominant multisystemic disorder caused by a microsatellite repeat expansion within intron 1 of the zinc finger protein 9 (ZNF9; also known as CNBP) gene. DM2 is generally characterized by a less severe phenotype than DM1 (myotonic dystrophy type 1). Despite a growing appreciation for the cardiovascular manifestations in DM1, cardiac involvement in DM2 has been less well characterized. In patients with DM2, cardiomyopathy has rarely been described. Case Presentation: This clinical vignette describes a rare case of DM2 associated cardiomyopathy in a 56-year-old male with DM2 who presented with palpitations, fatigue, reduced exercise tolerance, atypical chest pain, and lower extremity edema. He also reported proximal progressive muscle weakness for the past 20 years. Genetic testing confirmed DM2 due to a ZFN9 mutation. Physical exam showed normal vital signs, no jugular venous distention or edema, clear lungs, regular heart rate and rhythm without murmurs, and mild proximal muscle weakness. He additionally had hypertension, dyslipidemia, prior tobacco use (15 pack-years), well-controlled

fibrin thrombi and extravasation of red blood cells consistent with warfarin necrosis.

With a de

conditions, seeing multiple subspecialists, and poor medical record keeping in the healthcare system. To help improve outcomes in patients with excessive

medications to undergo a series of comprehensive medication reviews with our pharmacist and their primary care provider (PCP) in an effort to maintain an accurate medication list.

Methods: Patients could be included in the study if they were on >10 medications between October 2020 and August 2021. A comprehensive medication review was completed by our clinic pharmacist and total number of medications and any medication discrepancies were noted. The patients then completed a clinic appointment with their primary care provider dedicated to medication list review and education. Medications were again reviewed via pharmacist phone visit at 1 month and 6 months following PCP visit and discrepancies and total number of medications were noted again. Frequency of medication discrepancies, common categories of medication discrepancies and change in total number of medications were compared over the series of follow ups.

Results: 45 patients are actively enrolled in this study. 51% of these patients did not complete their dedicated PCP visit immediately following their initial pharmacist visit, 57% of them due to needing an acute visit (clinic, ED or

	lung and was initially treated with nivolumab (PD-1 inhibitor) and later impilimumab (CTLA-4 inhibitor). He developed autoimmune cutaneous toxicity so impilimumab was stopped. Nivolumab was continued for an additional 11 cycles before he was hospitalized for ongoing nausea, vomiting, and a 40 lb weight loss. The second case was a man with merkel cell carcinoma admitted to the hospital with fatigue, anorexia, and nausea with vomiting after receiving 7 cycles of pembrolizumab (PD_1 inhibitor). In both cases laboratory evaluation was consistent with secondary adrenal insufficiency and hypophysitis from CPI therapy. The CPI therapy was stopped. Both patients were started on glucocorticoid replacement therapy with complete resolution of symptoms at follow up visits. Conclusion: These cases demonstrate an increasingly common development of irAEs associated with CPIs. Given the growing number of patients treated with CPIs, providers will need to be aware of potential irAEs such as adrenal insufficiency which can be severe and potentially life threatening.
Reid Eggleston Dr. Thomas Hartman Dr. Laura Walkoff Dr. Eunhee S Yi Dr. Jay H Ryu Dr. Misbah Baqir	*Pulmonary ATTR Amyloidosis: Clinical, Radiologic and Pathologic Features and Outcomes* Introduction: Transthyretin amyloidosis is a disorder characterized by deposition of a misfolded conformation of the transport protein transthyretin (ATTR) most commonly in cardiac and nervous tissue causing clinical disease. Pulmonary amyloidosis, or deposition of ATTR in lung tissue, is a poorly characterized manifestation of this disease. We present the clinical course, imaging characteristics, pathology results, and outcomes of a cohort of patients diagnosed with pulmonary amyloidosis at an academic institution. Methods: We performed a retrospective descriptive review of 28 patients at Mayo Clinic between 2005 and 2020 with pulmonary ATTR amyloidosis (89% male, age at diagnosis 50- demographics, subjective symptoms, tissue biopsy results, and pulmonary function testing, imaging, and treatment data. Results: Patients were typically diagnosed after persistent dyspnea and abnormal chest imaging resulted in bronchoscopy yielding ATTR amyloidosis on biopsy. Most had a pre-existing diagnosis of cardiac ATTR amyloidosis. A similar number of patients had wild-type and hereditary disease. Patients typically had mild restrictive lung dysfunction with moderate reduction in diffusion capacity for carbon monoxide. Ninety-three percent had chest computed tomography imaging most commonly showing diffuse nodularity. Several patients had pulmonary cysts, pleural effusions, or noncalcified lymphadenopathy. Almost all patients had pulmonary vascular involvement and half had interstitial involvement on tissue biopsy. One-third received either anti-amyloid pharmacotherapy or organ transplant. Half of patients died by the time of study inclusion [Jan. 1, 2005-Dec. 31, 2020]. Conclusion: Pulmonary disease is a less common, but clinically important manifestation of ATTR amyloidosis. While there is heterogeneity in the presentation of these patients, there is a common symptomatology and disease course. Anti-amyloid therapies are a promising development with associated mortality benefit.
Benjamin Krehbiel Dr. Adam Wolfe Dr. David Tierney	*Outcomes and Clinical Characteristics of COVID-19 in Patients with Tuberculosis* Introduction: COVID-

certain comorbidities, such as increased age, diabetes, cardiovascular disease, and chronic obstructive pulmonary disease. Tuberculosis (TB) disease severity and outcomes correlate with similar comorbidities. Literature on COVID-19 infection in patients with TB is limited. This study evaluated the clinical characteristics, illness severity, and mortality of COVID-19 infection in patients with TB compared to patients without TB.

	he denied any symptoms of polydipsia, polyuria, vision changes or weight changes. He was adamant that his fasting glucose was 100-130 at home, which did not correlate with the elevated A1c. A fructosamine was therefore obtained which was mildly elevated at 312, correlating to an A1c of roughly 7.5%. As the A1c and fructosamine were discordant, the conclusion was the A1c was an unreliable test in this patient. An initial workup to evaluate for cause of falsely elevated A1c included thorough medication review and social history, lipid panel, BMP and CBC. As the workup proceeds, the patient is being evaluated for possible continuous glucose monitor and will have fructosamine testing done in the future instead of A1c. Conclusion: The A1c is often a reliable test, but this case demonstrates the potential imperfection of the test. The A1c relies on a non-enzymatic reaction between hemoglobin and glucose, so there are many factors that may impact hemoglobin and therefore A1c accuracy. These factors include inherited thalassemias, acquired anemias, certain supplements/medications, renal disease and several other causes. Although false A1c is relatively uncommon, it may be underdiagnosed. It is therefore important to correlate A1c with fasting glucose measurements and use alternative laboratory biomarkers as another measurement if there are any underlying disease processes that may interfere with A1c accuracy.
David Jonason Dr. Guru Trikudanathan **2021 MN-ACP Resident-Clinical Vignette finalist**	*A Case of MALT Lymphoma Masked as Gastric Varices Confirmed by EUS* Introduction: Gastric varices (GV) can cause gastrointestinal bleeding and increase morbidity and mortality in patients with cirrhosis. Upper endoscopy with direct injection of tissue adhesive glue or sclerosing agents is commonly utilized whenever there is no endoscopic ultrasound (EUS) expertise. However, alternative diagnoses can masquerade the endoscopic appearance and presentation of GV leading to suboptimal management and delays in diagnosis. We present a case of MALT lymphoma masked as GV on endoscopy and later correctly identified by Doppler EUS. This case highlights the importance of EUS in the evaluation of GV to aid appropriate management. Case Presentation: A 60-year-old male with a history of alcohol and NSAID use was admitted following a pre-syncopal episode. Physical exam was significant for tachycardia and hypotension. Labs were significant for a hemoglobin of 6.9 g/dL. Upper endoscopy showed isolated large gastric varices with stigmata of recent bleeding. However, there were no signs of cirrhosis with portal hypertension or gastric varices on CT scan. He was referred for EUS-guided coil/glue embolization during which he was found to have a large subepithelial lesion (SEL) in the gastric fundus. Doppler interrogation excluded underlying varix but revealed a large intramural hypoechoic mass. Two smaller SELs were also found, one with a deep ulceration and a visible vessel. Hemostasis was achieved with injection of Epinephrine and placement of hemostatic clips. Ulcers on biopsy with immunohistochemistry showed neoplastic CD 20 positive cells consistent with low grade B-cell lymphoma such as mucosa-associated lymphoid tissue (MALT) lymphoma. H.pylori was negative. He was subsequently referred to oncology for radiation treatment. Discussion: EUS exploration should be offered in cases where direct visual diagnosis of gastric varices with endoscopy is unclear or when there is no evidence of portal hypertension on cross-sectional imaging or in the absence of cirrhosis to exclude alternative diagnosis. EUS can distinguish varices

	located in the submucosa from other causes for prominent gastric folds. In this case, EUS aided in making a timely diagnosis of gastric SEL masquerading as varices which further impacted management. EUS should be preferred over conventional endoscopy for GV as it offers many conceptual diagnostic and therapeutic advantages including real-time imaging that enables detection and precise targeting of perforating and collateral vessels which often accounts for uncontrolled or recurrent bleeding.
Stephanie Kuhlman **2021 MN-ACP Resident-Clinical Vignette finalist**	*Severe Methemoglobinemia After Sodium Nitrite Ingestion* Introduction: 39-year-old female presented after an intentional sodium nitrite ingestion in an attempt to kill herself. Case Description: A 39-year-old female with a history of major depressive disorder, post-traumatic stress disorder, and borderline personality disorder presented to the ED by EMS after a suicide attempt. She reported intentionally ingesting 4 tablespoons of sodium nitrite and quickly after that called 911. She arrived to the ED about 20 minutes after the initial reported but was sitting up talking to him, so he rushed her to the ED. On arrival she was briefly awake and speaking but became lethargic and then began to have a seizure. She received lorazepam and Keppra and was intubated for airway protection. She initially received activated charcoal for the ingestion. Her methemoglobin level came back at 73 so she received methylene blue (2 doses total). Her skin was bluish gray on presentation and when the ED physicians placed an arterial line her blood was dark brown. Her blood pressures began to drop and she required at one point up to 3 pressors to maintain adequate blood pressure. Her lactate was 11.7, pH 7.15, and bicarb 11 initially. The SpO2 persistently read in the 60s. In the ED after receiving methylene blue, her methemoglobin level was improved to 29. She was admitted to the ICU and remained intubated. When sedation was stopped, she began to have occasional and focal myoclonus, so propofol was restarted. This developed into full body myoclonus vs seizure activity. She was started on a versed drip at that time with concern for seizure. An EEG was obtained that did not show any seizure activity. About 36 hours after admission she was weaned from sedation and able to follow commands. She was extubated successfully. She continues to have suicidal ideation and is pending admission to inpatient psychiatry. She does not appear to have significant neurologic damage. Discussion: A methemoglobin level reports the percentage of hemoglobin that has been oxidized to the ferric state and cannot carry oxygen. She presented with an initial level of 73% which means only 27% of the hemoglobin in her body was capable of carrying oxygen. With this high of a level it is interesting that she did not have any kidney or liver damage on admission. She appeared to initially have hypoxic brain injury with the myoclonus and non-responsiveness of of sedation, although this now appears to be a toxic effect of the sodium nitrate more so than a hypoxic injury as she demonstrated good neurologic recovery.
Nermine Abdelwahab Dr. Emily Ewan Dr. Kristin Inman	*That Blood Pressure, You Gotta be Kidney-ing Me!* Case Presentation: A 50 year old female with history of bilateral renal cell carcinoma (RCC) s/p partial left nephrectomy, ESRD on hemodialysis, and hypertension was admitted for hypertensive emergency with a blood pressure

	diagnosis can be challenging and a broad differential of headache in patients who present to ambulatory clinic is essential. Classic signs and symptoms (pain, Horner syndrome, cranial nerve palsy, cerebral ischemia) or a high index of suspicion as in this case should prompt angiography for diagnosis. The management of carotid artery dissection is focused on preventing thromboembolic complication. Our patient was treated with daily aspirin 325mg and clopidogrel 75mg with a follow up MR angiogram in three months to determine duration of therapy.
Momen Alsayed	*I Broke my Heart in Connecticut; A Case of Lyme Carditis with High Degree AV Block* Introduction: Lyme disease is the most common vector-borne disease in the US. It is caused by the bacterium Borrelia burgdorferi and rarely, Borrelia Mayonii which is transmitted to humans through black-legged ticks (1). In 2019 alone, there were 23,453 per 100,000 confirmed cases in the US (2). Lyme carditis remains an uncommon complication in the early disseminated phase of the disease. In this report, we hope to shine a light on the importance of cardiac workup in this population. Case Presentation: A 41-year-old female with a past medical history of class 3 obesity, OSA, HTN, PCOS, saphenous vein thrombosis, and splenectomy post-MVA, presented to the clinic feeling fatigued for 1 month after returning from a Cabin trip to Connecticut. A week before presentation the patient had low-grade fevers, a disseminated rash on her abdomen, and low home blood pressure and heart rate readings. Initially, she denied palpitations, chest pain, shortness of breath, lightheadedness, or syncope. Her heart rate was in the lower 50s bpm but her vitals were otherwise normal. Erythema migrans was noted on the upper abdomen, but the exam was largely unrevealing. Her initial EKG showed evidence of intermittent high degree AV block. Initial workup was negative for COVID-19, electrolytes and hepatic derangements, and thyroid dysfunction. Her WBC count was 11.36k/cmm and her neutrophils 7.99k/cmm. During admission, serial EKGs revealed progression in the degree of her AV block to persistent high degree AV block and evidence of intermittent ventricular escape rhythm. TTE showed borderline increased LV wall thickness and asynchronous septal-apical motion but was negative for structural, valvular, or wall motion abnormalities. A temporary cardiac pacemaker was placed and the patient was started on 3g IV ceftriaxone after a presumptive diagnosis of Lyme carditis was made. On the 3rd day, Lyme IgG & IgM western blots were positive and the patient was continued on IV ceftriaxone. After 7 days of antibioticcs, the patient was reassessed and the pacemaker was removed after sustained improvement and resolution of high degree AV block on repeat EKGs. She was switched to 100mg oral doxycycline BID on discharge for a total of 21 days of antibiotics. She was then seen by cardiology 1 day after completing her antibiotic course, was asymptomatic and her EKG was normal with complete resolution of AV block. Conclusion: This case highlights the importance of early recognition and management of Lyme disease to prevent cardiac complications such as acute AV nodal disorders which can rapidly evolve to a high degree AV block and lead to cardiac morbidity and mortality. Having a higher clinical suspicion for Lyme carditis, especially in patients with travel history to endemic areas who present with bradycardia or AV block with or without cardiac symptoms,

	allows for timely management which includes temporary cardiac pacing and antibiotics to prevent negative outcomes. 1. (CDC,2021), https://www.cdc.gov/lyme/index.html 2. (CDC, 2019), https://wonder.cdc.gov/nndss/static/2019/annual/2019-table2j.html
Hadiyah Audil Dr. Jennifer Duke Dr. Patricio Escalante	*Pulmonary-Cutaneous Fistula Associated with Empyema Necessitans due to M. Avium Complex* Case Presentation: A 67-year-old woman with a history of right breast cancer status post chemoradiation, recent longstanding right-sided pneumonia (diagnosed at outside facility, unsuccessfully treated for months with oral antibiotics), and BMI 13 presented with acute development of a purulent-draining cutaneous fistula on her right chest wall. CT chest showed a right pulmonary-cutaneous fistula (PCF), pleural effusion, and pneumothorax. An occlusive dressing was placed over the fistula and she was started on vancomycin and piperacillin-tazobactam. Thoracic Surgery was consulted and recommended creation of an open thoracic window, as her malnutrition precluded muscle flap creation; the patient declined this in favor of drain placement. Cultures of the drainage grew Mycobacterium avium complex (MAC), and she was started on rifampin, ethambutol, azithromycin, and amikacin with outpatient follow up. Discussion: To our knowledge, PCFs have not been previously reported with MAC. Risk factors for PCF development include prior thoracic surgery/procedures; thoracic radiation; malignancy; malnutrition; and empyema necessitans associated with chronic lung infections (particularly with M. tuberculosis, M. abscessus complex, and Actinomyces), wherein extension of the infected pleural space to the chest wall causes skin/soft tissue erosion. Patients should initially be placed on airborne precautions given the association with tuberculosis. CT chest should be obtained to evaluate for co-existing intrathoracic pathologies. Thoracic Surgery should usually be consulted for consideration of surgical treatment. If drainage is observed, bacterial/fungal cultures should be obtained, and broad-spectrum antibiotics should empirically be started. Although morbidity and mortality can be high for untreated PCF, prognosis is usually good with appropriate management.
Connor Buechler Dr. Tom Freeman	*A Tale of Two Blisters* Introduction: Systemic Lupus Erythematosus (SLE) is a chronic autoimmune connective tissue disorder of unknown origin with wide-ranging clinical manifestations. In the absence of widely accepted diagnostic criteria, diagnosis is generally guided by classification criteria, recognition of classic features, and exclusion of alternative diagnoses. Porphyria cutanea tarda (PCT) is a metabolic disorder caused by altered activity of the heme biosynthetic enzyme uroporphyrinogen decarboxylase (UROD) in the liver, and is much rarer in clinical practice than SLE. Blistering skin lesions are the predominant clinical manifestation of PCT, and over 80% of cases are due to an acquired deficiency rather than inborn mutation. Case Presentation: A 64 year old male with a past medical history of coronary artery disease presented with a rash beginning on his trunk one month prior to presentation. A dermatology visit at that time noted a widespread annular erythematous and edematou

interface dermatitis with apoptotic basal keratinocytes, moderately dense superficial perivascular lymphocytic infiltrate, and no alteration of dermal collagen. The patient was placed on topical triamcinolone cream which proved ineffective. The patient presented to the ED three weeks later with progressive desquamation and mucositis prior to ultimately being transferred to our hospital with concern for toxic epidermal necrolysis. On admission, he had a severe erythematous blistering and erosive rash with occasional tense

	requiring eventual ICD placement, recurrent pulmonary edema and progressive renal failure despite maximal therapies and initial response to chemotherapy. Ultimately discharged with family on hospice cares on hospital day 27. Conclusion: The relatively poor prognosis of primary AL amyloidosis compared to other fibril types is largely related to the frequency of cardiac involvement, including a higher incidence of LVH and intracardiac thrombosis [1]. Unfortunately atypical systemic symptoms can be presenting features of AL amyloidosis progression. Early goals of care discussions and involvement of palliative medicine should be considered given higher risk for sudden fatality. References: [1] Choukair MO, Halawi A, Nehmeh A, Kazma H. Role of Echocardiography in the Diagnosis of Light Chain Amyloidosis: A Case Report and Review of Literature. Cureus. 2020;12(11):e11377. Published 2020 Nov 8. doi:10.7759/cureus.11377 [2] Kastritis E, Dimopoulos MA. Recent advances in the management of AL Amyloidosis. Br J Haematol. 2016 Jan;172(2):170-86. doi: 10.1111/bjh.13805
Karan Chohan Dr. N. Nora Bennani	*A Case of Cartilage-Hair Hypoplasia and Non-Hodgkin Lymphoma* Introduction: Cartilage-hair hypoplasia (CHH) is a rare autosomal recessive chondrodysplasia often characterized by short-limbed short stature, hypoplastic hair, combined iCon

fibrosis and has been found to be quite steroid responsive, however no second line treatment has yet been established.

Case Presentation: We present a case of a 77-year-old woman with a history

with unexplained macrocytic anemia noted during routine monitoring labs. She was seen by her rheumatologist for her autoimmune arthritis which had been well controlled with methotrexate, adalimumab, and as needed ibuprofen for many years. Complete blood count was ordered for monitoring while on methotrexate and she was found to have precipitous drop in her hemoglobin over the past three months. Gastrointestinal loss in the setting of ibuprofen use was first considered, however the patient refused endoscopic workup and was eventually referred to hematology. At that time, she had elevated inflammatory markers with a significantly elevated sedimentation rate and c-reactive protein. In addition, she was noted to have an elevated ferritin and iron studies which showed a mixed picture. Because of this, a broad workup was ordered including a bone marrow biopsy. The bone marrow biopsy results showed a hypercellular marrow with moderate reticulin fibrosis consistent with secondary autoimmune myelofibrosis associated with rheumatoid arthritis. After obtaining these results she was started on high dose prednisone for treatment.
Unfortunately, the patient subsequently developed a pseudomonas septic arthritis of her sacroiliac joint and required prolonged treatment with intravenous antibiotics, along with discontinuation of high dose prednisone and adalimumab due to the infection. She was thus started on treatment with rituximab after discussion between her hematologist and rheumatologist. This did eventually lead to a modest improvement of her hemoglobin level.

Discussion: Autoimmune myelofibrosis is a clinically rare disease that must be differentiated from other causes of myelofibrosis. A clonal malignant disease is seen in patients who have gain of function mutations resulting in disease such as janus kinase 2 (JAK2) or calreticulin mutation (CALR). Autoimmune myelofibrosis can be either secondaryJETJET0 G[()-4(t)6ETybo0

	following 24 hours, she developed ophthalmoplegia, limb ataxia, bilateral nystagmus, urinary retention, and tremors. Initial differential was highly concerning for pontine osmotic demyelination syndrome and imaging was ordered to evaluate. MRI demonstrated a nonspecific T2 hyperintensity of the medial hypothalamus and periaqueductal gray, and spinal MRI demonstrated an abnormal diffuse signal edema. Neurology was consulted overnight due to the new neurologic findings and this abnormal imaging. Osmotic demyelination was not consistent with her imaging and was ruled out. A lumbar puncture was performed by neurology and the cerebral spinal fluid was remarkable for white blood cell count of 161, and elevated IgG. Given the absence of infectious symptoms, there grew an increasing suspicion for an underlying rheumatologic process. Approximately one week into admission she developed cranial nerve VII palsy, and progressive lower extremity weakness. Relevant serology studies were positive for ANA, anti-Ro, and anti-aquaporin-4. History, imaging, and serology findings solidified a diagnosis of neuromyelitis optica spectrum hyponatremia was secondary to a combination of hypovolemia as well as SIADH caused by her NMOSD. Immediately following diagnosis, she was started on plasmapheresis and 1g daily of methylprednisolone for five days, followed by a prolonged prednisone taper over at least six months. She received a Rituximab infusion, with plans to undergo infusions every six months as an outpatient. The patient was ultimately discharged to an acute rehabilitation facility after a four-week admission. Since initiation of treatment she has had moderate improvement of upper extremity weakness, however continues to suffer from profound lower extremity weakness, neurogenic bladder, and neuropathic pain. Discussion: Though atypical, intractable nausea and vomiting can be the only presenting symptoms of NMOSD. This progressive disease is characterized by periods of relapse leading to cumulative disability, which can be fatal. It is critical to recognize this presentation, given that earlier initiation of immunotherapy has proven to decrease rate and frequency of relapse, and improve long-term outcomes. Therefore, it is important to remember that not
Jissy Cyriac	, ¶ P %DFNHG 8S 5HQDO)DLOXUH DQG 0\RF *Bladder Incarceration* Background: Obstructive uropathy can lead to urinary retention and acute kidney injury. We present a case of obstructive uropathy secondary to inguinal hernia with bladder incarceration leading to renal failure and myocardial infarction. Case Presentation: An 80-year-

bilateral hydronephrosis. Labs were notable for elevations in creatinine (14.83

dilemma for internists due to its infrequent clinical presentation and wide differential diagnosis that includes allergic, infectious, rheumatologic, neoplastic, and less commonly idiopathic disorders. Eosinophilia is defined as

and hypereosinophilic syndrome (hypereosinophilia plus end-organ dysfunction). We present a case of hypereosinophilic syndrome with pulmonary and renal dysfunction.

Case Presentation: A 76-year-old Somali-immigrant was admitted to an outside hospital with several weeks of progressive dyspnea and wheezing. His past medical history was significant for ischemic cardiomyopathy, type 2 diabetes mellitus, and chronic kidney disease. His social history was significant for having lived in Somalia, Yemen, and Indonesia prior to immigrating to Minnesota five years earlier. At the outside hospital, he was treated for community-acquired pneumonia without improvement. Due to his

glucocorticoids and transferred to our hospital for further evaluation.

During his hospitalization, a broad evaluation was obtained to identify the etiology of his hypereosinophilia. His evaluation was significant for a peak

(normal <3%), elevated IgE to 817 kU/L (normal <214), positive toxocara antibodies, and peak creatinine 2.66 mg/dL (baseline creatinine 1.3-1.4). The remainder of his allergic, infectious, rheumatologic, and hematologic evaluation was unrevealing. He was treated with albendazole for toxocariasis and inhaled corticosteroids for his wheezing. His eosinophilia downtrended and clinical symptoms resolved. Bone marrow biopsy was deferred due to his clinical improvement.

Discussion: We present a case of hypereosinophilic syndrome secondary to toxocariasis. The clinical approach to peripheral eosinophilia should begin with a broad multisystem evaluation to identify the underlying etiology. Clinicians should also evaluate for evidence of end-organ involvement bearing in mind hypereosinophilic syndrome most commonly involves the pulmonary, cardiac, renal, gastrointestinal, central nervous, and/or cutaneous systems.

	hospitalization, while CT and MRI studies of the cervical spine did not clearly demonstrate infection, he had been found to have MSSA bacteremia and leukocytosis of 38 x 109/L prompting osteomyelitis concerns. The patient had also developed left lower extremity swelling, with negative DVT ultrasound, precipitously dropped from 15.5 to 7.9 secondary to a gastrointestinal bleed which was treated supportively. He was transferred due to concern for higher level of care. After transfer, the patient continued to have significant neck tenderness and lower extremity pain. He was treated with IV cefazolin for the MSSA bacteremia. Reexamination of the prior CT and MRI imaging of the cervical spine demonstrated evidence of unattached anterior cervical hardware from the spine. The patient continued to have significant hemoglobin drop, prompting investigation with EGD and colonoscopy. EGD provided direct visualization of his cervical hardware eroding into the upper third of the esophagus, with follow-up esophagram negative for extravasation around this hardware erosion. TTE showed no evidence of endocarditis or infected thrombi. He was planned for complex esophageal repair and anterior cervical plate removal. Discussion: For patients with indwelling hardware, malfunction always remains on the differential. For this patient, his chronic inflammatory state and prednisone use may have contributed to migration of spinal hardware. While his GI bleed resulted in escalated care, more subtle symptoms of this erosion were present such as progressive dysphagia and more recent neck pain. It is important to evaluate indwelling devices and hardware, especially in patients with localized symptoms.
Aoife Feighery	*The Cecal Bascule: A rare form of large bowel obstruction* Introduction: A cecal bascule is a rare form of bowel obstruction where the cecum folds upon itself in an anterior-caudal direction along the ascending colon. Cecal bascules account for only 0.05-0.4% of large bowel obstructions. In contrast to other forms of mechanical obstruction, the cecum in a bascule folds along a transverse line, without torsion. This movement is likened to a see- experienced in other forms of bowel obstruction: abdominal pain, distension, constipation, and occasionally respiratory distress due to abdominal hypertension. Diagnosis usually involves an abdominal Xray showing a dilated cecum in the right upper quadrant without the axial twisting or the be utilized for further characterization and identification of perforation or ischemia. Initial management incudes colonic decompression, pain control, and respiratory support. Ultimately, the treatment of a cecal bascule is surgical, typically right hemicolectomy. Case Presentation: An 80-year-old male with a history of COPD, congestive heart failure, paroxysmal atrial fibrillation, and chronic kidney disease

	for impending bowel perforation. Upon arrival to the ICU, the patient required bilevel positive pressure ventilation. His abdomen was firm, distended, and markedly tender in the right upper quadrant. Bowel sounds were diminished. CT showed a dilated cecum flipped into the right upper quadrant, consistent with a cecal bascule. There was no free air, no upstream bowel dilation, and no cecal wall thickening or mesenteric edema to suggest acute volvulus. A nasogastric tube was placed for decompression. An aggressive bowel regimen was initiated without effect. Hypaque enema and endoscopic colonic detorsion were unsuccessful. After exhausting non-operative options, the patient was taken for explorative laparotomy, which confirmed a cecal bascule with impeding rupture. Right hemicolectomy with end ileostomy was performed. Conclusion: Bowel obstruction is common within both inpatient and outpatient settings. Typically, non-mechanical or functional obstruction is managed medically, while mechanical obstruction, such as colonic mass, volvulus, and intussusception requires surgical intervention. Cecal bascule is an uncommon form of mechanical obstruction not responsive to aggressive medical or endoscopic therapy. It is paramount for physicians to identify such cases early in the clinical course and to involve a surgical team for definitive intervention to prevent intestinal ischemia, necrosis, or perforation.
Nouran Felo Dr. Siri Urquhart Ddr. Sumedh Hoskote	*When the Patient Reads the Textbook ±A Classic Case of Thrombotic Thrombocytopenic Purpura* Introduction: Thrombotic thrombocytopenic purpura (TTP) is a thrombotic microangiopathy characterized by thrombocytopenia, microangiopathic hemolytic anemia (MAHA), and organ ischemia; a medical emergency that should be recognized quickly to initiate prompt medical treatment. Although ADAMTS13 activity, which can confirm the diagnosis of TTP takes time to result, TTP can be recognized clhe

inhibitor, confirming his diagnosis. He was continued on daily PLEX until platelet count recovered and was started on high-

	Mexiletine and Amiodarone. An ICD was not placed as per patient preference. For treatment of the giant cell myocarditis, she was started on pulse-dose IV steroids followed by high-dose oral steroids with a prolonged taper. Cyclosporine and Azathioprine were also initiated for immunosuppression. She was discharged in improved condition with appropriate cardiology follow-up. Conclusion: Though giant cell myocarditis is rare, its diagnosis is important as the treatment differs from other forms of myocarditis. For giant cell myocarditis, prompt treatment with a combination of immunosuppressive medications provides improved prognosis. Even with treatment, the risk of ventricular arrhythmias remains elevated. Ultimately, cardiac transplantation is often warranted, though recurrence is common.
Vanessa Gow-Lee Dr. Elie Berbari	*EBV Meningitis as Manifestation of Infectious Mononucleosis* Introduction: Although Epstein-Barr virus in the central nervous system has been well-characterized in immunocompromised patients, CNS involvement in immunocompetent hosts with primary infection is not. Furthermore, the significance of detecting EBV by PCR in the cerebrospinal fluid is not always clear, especially in immunocompetent patients. EBV has been described to cause a variety of CNS syndromes, including meningitis, encephalitis, cerebellar ataxia, acute disseminated encephalomyelitis, and myelitis. We describe here a case of mononucleosis in an immunocompetent adult who presented with aseptic meningitis which was attributed to EBV. Case Presentation: Our patient was a 27 year-old woman with minimal comorbidities (ADHD, chronic sinusitis) who presented with several weeks of fatigue, myalgias, left-sided abdominal pain, and a severe occipital and retro-orbital headache with associated neck stiffness, photo- and phonophobia. She denied any rash, pharyngitis, any new romantic/sexual partners, or high-risk sexual activity. She had nuchal rigidity on exam, but no lymphadenopathy or other focal neurological deficits. Laboratory tests showed a lymphocytic leukocytosis (WBC 14.6 x 10^9 cells/L with >80% lymphocytes, with smear showing atypical lymphocytes) with a new transaminitis and hyperbilirubinemia (AST 324 U/L, ALT 387 U/L, tBili 3.8 mg/dL, dBili 2.8 mg/dL). EBV serologies revealed positive IgM and negative IgG, indicating acute infection. Abdominal imaging showed splenomegaly of 17.2 cm. MRI of the brain was normal without parenchymal changes or hydrocephalus. LP revealed an elevated opening pressure (>40 cm H2O) and lymphocytic pleocytosis (75 cells with 81% lymphocytes), slightly elevated protein (66 mg/dL), and normal glucose (52 mg/dL). Gram stain and culture were negative and a broad-range PCR panel was positive only for EBV. She was initially treated conservat



Alexand

lungs and axillary lymph nodes. Biopsy revealed noniopsy revea

showed atypical lymphocytes.

	squamous cell carcinoma. He then underwent bronchoscopy with bronchoalveolar lavage (BAL) with the immunocompromised diagnostic panel. BAL returned positive for pneumocystis pneumonia and he was transitioned from azithromycin to high dose IV trimethoprim sulfamethoxazole and prednisone 80 mg daily with plans for total of 3 weeks of corticosteroids. The etiology of his immunocompromise was unclear but initial thought was given to recurrence of his squamous cell carcinoma, rheumatoid arthritis (or a concomitant inflammatory condition), long-term methotrexate use, or hematologic malignancy. His initial work up demonstrated anemia and thrombocytopenia. He underwent bone marrow biopsy which demonstrated a new large B cell lymphoma which was the most likely source of immunocompromise and contributed to his dyspnea, fatigue, and weight loss. After learning of this diagnosis, he elected to pursue comfort measures and was discharged to hospice, where he passed away surrounded by his family. Conclusion: This case demonstrates several key lessons. First, it is important to maintain a broad differential diagnosis for dyspnea even during the midst of a pandemic where recency bias (a cognitive bias that favors recent diagnoses over historic ones) can cloud clinical judgement. Second, any patient diagnosed with PCP pneumonia needs to undergo evaluation for underlying immunocompromise, with focus on recent steroid administration, hematologic malignancy, underlying inflammatory conditions, or other solid malignancies if the patient has tested negative for HIV. Lastly, it has been demonstrated that PCP in patients without HIV is more difficult to treat and carries a higher mortality.
Jeffrey Heller Tanisha Ronnie	*Rheum for Improvement - A Case of Transverse Myelitis and the Need for Early Rheumatologic Workup* Introduction: Transverse myelitis is characterized by weakness, sensory loss, pain, and bowel and bladder dysfunction secondary to an inflammatory lesion of the spinal cord. It is a common initial presentation of systemic lupus erythematosus (SLE). In fact in 50-60% of patients with SLE and transverse myelitis, transverse myelitis is the initial presentation, prior to other more common sequelae of the disease (1,2). Thus, diagnostic evaluation for rheumatological causes should be pursued when transverse myelitis is suspected. Case Presentation: An otherwise healthy 31 year-old woman, with no significant past medical history, presented to the hospital with three weeks of worsening nausea and vomiting. On day 3 of her hospitalization, the patient developed severe weakness in her lower extremities, urinary retention, and bowel incontinence. MRI showed diffuse inflammation in the spinal cord, suggesting extensive longitudinal myelitis, defined as transverse myelitis involving three or more contiguous vertebral segments of the spinal cord. PLEX therapy and IV steroids were initiated on day 6, given the unknown etiology of spinal inflammation. On hospital day 8, there was no significant improvement in her symptoms with CBC notable for worsening pancytopenia. With the transition of the medical team, further history taking revealed a 6-month history of joint pain and eye inflammation. The patient was not successfully diagnosed with SLE until hospital day 10, and treatment was started on hospital day 12, approximately one week after the development of extensive longitudinal myelitiinence. Mer symptEW08.74 108I5ETQq48 reu 11

Conclusion: This case highlights the importance of thorough history taking and pursuing a rheumatologic workup in patients presenting with transverse myelitis. When considering the differential for transverse myelitis there are a number of etiologies that must be considered including vascular lesions, structural lesions, postinfectious, post-vaccination, paraneoplastic, autoimmune, demyelinating and metabolic causes (3, 4). A detailed history with questions asking about visual changes, constitutional symptoms, dermatologic findings, joint and muscle pain, history of pregnancy loss, and overall timing and onset of symptoms can guide the workup. Imaging is the first step in the workup and is twofold in its purpose. It provides radiologic confirmation of the diagnosis and can also indicate whether a compressive lesion is present. Spinal MR.74 rannpb163.58 550.3 356.37tms c 9(t)-he ofm(e)

replaced with illicit analogs of fentanyl, which are highly lipophilic, distribute widely to adipose tissue, and lead to protracted excretion. This pharmacokinetic profile increases the risk of precipitated withdrawal after starting buprenorphine. Microinduction, also known as microdosing, is an effective method for starting buprenorphine in the hospital setting that decreases the risk of precipitated withdrawal while offering acute pain control with other opioids. With improved symptom control, patients with OUD are less likely to leave against medical advice, leading to improved clinical outcomes.

	the patient was transferred to an outside hospital for surgical VSD repair, aneurysmectomy, and CABG. The patient passed away two weeks later. Discussion: This case illustrates a typical course for the most common post-MI mechanical complication: rupture of the IS. Because infarcted myocardium is not amenable to surgical repair, the prognosis is poor even with prompt identification. However, early diagnosis can help expedite clinical decisions and improve the chances of a good outcome. A large majority of post-MI septal ruptures result in a characteristic holosystolic murmur, and therefore diligent cardiac auscultation can result in diagnosis prior to the onset of cardiogenic shock and associated complications.
Isla Johnson Dr. Gretchen Colbenson	*Taming of the Sprue: Celiac Disease-Associated Neutropenia as Cause of Prolonged Neutropenia* Introduction: Neutropenia is a hematologic abnormality with a broad differential diagnosis, and especially in hospitalized patients, can often be attributable to illness or medication effect. However, prolonged neutropenia merits extensive workup as it confers increased risk of infection and, with Physicians should be aware of uncommon etiologies, such as Celiac Disease (CD), during workup for neutropenia. Case Presentation:

infection (Vijayvargiya et al., Blood 2015). In the case of severe infection, G-

viral panel and COIVD test were negative. Basic labs were unremarkable. Chest X-ray was negative, CTPA was unremarkable except for mildly enlarged left hilar lymph nodes most likely reactive. ECG showed nonspecific subtle diffuse ST-segment abnormalities with normal sinus rhythm, exercises stress echo was negative for ischemia or pericardial effusion. The patient did not receive antibiotics and improved with NSAIDs. Impression was a mild viral illness, and he was discharged with a plan of near follow-up.

On 4/18 presented again to our ER with persistent fever (reaching 102.5),

	Pulmonary HTN, A-fib/A-flutter (was on amiodarone 200 mg daily for 4 months and was then stopped 1 month before presentation), CAD, CKD, T2DM, and HTN. His home nurse saw him and noted an O2 sat in the 80s, usually requiring 1-2L but now requiring 4L O2. He presented to the ED reporting acute on chronic SOB. The patient was tachypneic, and normotensive. Initial labs with BMP significant for Cr 2.3 mg/dl. VBG significant for pH 7.42, CO2 of 47 mmHg, Bicarb of 30 mEq/L. CBC showing Hgb of 6.7 g/dl. NT proBNP of 1720 pg/mL, with unremarkable troponins. PT of 35.2 sec and INR of 3.0. Initial imaging was consistent with pulmonary edema. The patient received 1 unit of RBCs and was started on aggressive diuresis with no response. He went from 4L O2 NC to requiring BiPAP and was transferred to CMIC for that reason. Cardiology and pulmonology were consulted. Additional workup included TTE, which was unremarkable. Follow-up CXRs with worsening opacities. ABG showed a pH of 7.28, CO2 of 81, and Bicarb of 35. Follow-up VBGs with persistently elevated CO2 ranging from 70s- hypoxia, and worsening CXRs were most likely due to amiodarone pulmonary toxicity. He was started on glucocorticoids, resulting in remarkable improvement in SOB. Discussion: Amiodarone pulmonary toxicity incidence ranges between 5-13%(2). Typically manifests as diffuse interstitial pneumonitis with varying degrees of fibrosis. Risk factors for APT include a high dose (>400 mg/day), duration > 2 months, increased age, and preexisting lung disease(1). Treatment is stopping amiodarone and initiating glucocorticoids. This case shows that while it does not have to be at the top of the differential diagnosis, it should still be relevant in patients exposed to amiodarone. Being a widely used medication, clinicians should be aware of the possibility that despite stopping amiodarone, its adverse effects may still present later, resulting in a need to consider targeted monitoring allowing for early diagnosis and management, minimizing morbidity and mortality. References: 1. Wolkove N, Baltzan M. Amiodarone pulmonary toxicity. Can Respir J. 2009;16(2):43-48. doi:10.1155/2009/282540 2. Schwaiblmair M, Berghaus T, Haeckel T, Wagner T, von Scheidt W. Amiodarone-induced pulmonary toxicity: an under-recognized and severe adverse effect? Clinical research in cardiology: official journal of the German Cardiac Society. 2010;99(11):693-700. doi:10.1007/s00392-010-0181-3
Jonathan Knott Dr. Caroline Matchett Dr. Matthew Koster	*Treatment-Resistant Autoinflammatory Disease: A Unique Case of VEXAS Syndrome in a 60-Year-Old Male* Introduction: VEXAS syndrome is a newly identified adult-onset inflammatory syndrome.1 It is characterized by a somatic mutation affecting methionine-41 in the UBA1 gene resulting in diminished cellular ubiquitylation and hyperinflammation affecting multiple organs, including the skin, lungs, blood vessels, and cartilage.1-2 VEXAS syndrome is associated with considerable morbidity and mortality as symptoms are refractory to typical treatments including steroids and steroid-sparing agents.2 Case Description: A 60-year-old male, non-smoker originally presented to an outside hospital with chest pain, left-sided spontaneous pneumothorax, and diffuse ground glass opacities (GGO). He then developed recurrent fever, thrombophlebitis, urticarial dermatosis, episcleritis and cytopenias with elevated acute phase reactants. Comprehensive infectious evaluation was

negative. Bone marrow was hypercellular but did not disclose overt malignancy. He responded to high-dose glucocorticoids, but symptoms returned on tapering below 20mg/day. Trials of dapsone and omalizumab were unsuccessful for control of his rash.

He presented to Mayo Clinic for evaluation with active symptoms while on mycophenolate 1000 mg twice daily and prednisone 20 mg. A repeat bone marrow revealed vacuolated erythroid and granulocytic precursors. He was hospitalized due to worsening dyspnea, fever and rash. Chest CT revealed progressive diffuse GGO. Bronchoscopy with broad infectious workup was again negative. Skin biopsy showed neutrophilic dermatosis. The constellation of symptoms was concerning for VEXAS and genetic testing

revealed a WBC count of 20,000 k/cmm and a lactate of 3.9 mmol/L.

Imaging studies including an EKG, CXR, and CT scans of the head, chest, abdomen, and pelvis were unremarkable except for scattered nodules in the right upper lobe of the lung on CT scan. Patient was treated empirically for possible community acquired pneumonia. Blood cultures resulted with methicillin sensitive Staphylococcus aureus and antibiotics were narrowed to

	10.9 cm by 7.3 cm, as well as a large pleural effusion and moderate pericardial effusion. Laboratory evaluation was notable for hemoglobin 10.1 g/dL and leukocytosis of 23.57/µL with 91% neutrophils. Alpha-fetoprotein (AFP) and beta-HCG were normal and lactate dehydrogenase was elevated. Cytology by CT-guided mediastinal biopsy was consistent with large B-cell lymphoma with extensive necrosis. Pleural fluid was positive for malignant cells. PET-CT scan showed FDG-avid large anterior mediastinal mass and mediastinal lymphadenopathy without other FDG-avid disease. Bone marrow biopsy and lumbar puncture were negative for malignancy. She met with reproductive endocrinology for fertility preservation prior to starting appropriate chemotherapy. Discussion: This cETQqc4o163.58 449.11 3nes the clinical presentation and dia evalion of an anterior mediastinal mass in a young female patient. Minimal response to antibiotics for pneumonia in young p9.11atients should prompt furth imaging. Appearance on high-resolution CT is important in differentiating etiology of AMM. This patient had a large hQqterogeneous mass with associated pleural effusion, suggesting the more aggressive large B-cell teratomas are diagnosed by imaging alone, and a homogeneous, lobulated mass raiQqs suspicion for thymoma. For the general internist evaluating an anterior mediastinal mass, demographic information, clinical symptoms, and appearance on imaging are critical for appropriate diagnosis and 9.11 3aging pri to oncology involvement.
Katy Lehenbauer	*Stress Cardiomyopathy Within the COVID-*

	possible hereditary cancer syndrome affecting both male and especially female relatives, further investigation into HDGC by obtaining prior medical records and slides, and/or CDH1 genetic testing, should be considered. There are screening and preventive treatment implications for the rest of the -minute upper endoscopy every 6-12 months with multiple random biopsies and biopsies of subtle lesions, beginning 5-10 years prior to the earliest cancer diagnosis in the family along with H. pylori screening, prophylactic total gastrectomy, as well as breast cancer screening and prophylactic surgery.
Rosemary Liu Dr. Rena Singleton	*REMAPping COVID-19* Introduction: Effective goals of care conversations have become increasingly important with the onset of the COVID-19 pandemic, particularly considering that COVID-19 can cause significant morbidity and mortality even in relatively healthy individuals. The REMAP framework provides guidance for how to navigate these complex conversations in a patient-centered way. Case Description: A previously healthy 80 year old male presented with four days of shortness of breath, dry cough, dizziness, fatigue, decreased appetite, and low-grade fever. After being found to be hypoxic to 85-80% on room air during a visit to his primary care doctor, he was sent to the ED where vitals were temperature of 100.7F, respiratory rate of 18, pulse of 89, blood pressure 121/77, and oxygen saturation of 78%. Physical exam revealed crackles in bilateral lower lung fields. CXR demonstrated bilateral interstitial changes concerning for COVID pneumonia, and COVID testing returned positive. Initial workup was notable for leukocytosis of 12.9, elevated AST (106), BNP (367), CRP (205), LDH (449), ferritin (823). Due to elevated D-dimer (1.59), CT chest was obtained which showed bilateral multilobar pulmonary space

	transitions in care are anticipated. It provides a template for conducting effective goals of care conversations that promote shared decision making with patients and their families. 1. Suresh, E., et al., Acute pancreatitis with pseudocyst formation in a patient with polyarteritis nodosa. J Rheumatol, 2005. 32(2): p. 386-8. 2. Flaherty J, Bradley EL 3rd. Acute pancreatitis as a complication of polyarteritis nodosa. Int J Pancreatol. 1999 Feb;25(1):53-7.

Caroline Matchett
Dr. Lillian Wang

	lung lesions was extensive with both unifying and separate diagnoses considered. With a history of positive toxoplasmosis IgG, compatible brain imaging findings, and toxoplasmosis being the most common cause of rim-enhancing lesions with a CD4 count <100, CNS/pulmonary toxoplasmosis appeared most likely. Mycobacterial infection, nocardiosis, bacteremia with septic emboli, fungal (i.e. Histoplasma, Cryptococcus, Pneumocystic jiroveci pneumonia (PCP)) or viral infections, lymphoma, or a combination of these were among other investigated diagnoses. He was empirically started on high-dose Bactrim and intravenous vancomycin and ceftriaxone, with the latter two discontinued after negative blood cultures and normal echocardiogram. Unfortunately, lumbar puncture was contraindicated due to edema, but serologies, cultures from sputum and bronchoalveolar lavage (BAL), and molecular tests were obtained. BAL was positive for P. jiroveci and toxoplasmosis. After two weeks on Bactrim, repeat brain MRI showed significant reduction in lesions and edema, while chest CT seemed to demonstrate progression of disease. Rare Candida albicans on sputum culture only was unlikely to be contributing to his lung disease, but nonetheless fluconazole was started for oral thrush. Testing for endemic mycoses, Aspergillus, Mycobacterial infections, and Echinoccocus was unremarkable. In consultation with pulmonary and radiology it was determined that the chest imaging was evolving in conjunction with treatment for presumed PCP, instead of worsening, and the patient remained on Bactrim at discharge after 5 weeks. His cognition slowly improved but he was still not at his baseline. Discussion: In patients with untreated HIV with new illness, it is important to consider possible opportunistic infections or malignancies based on CD4 count. This patient with a CD4 count <100 would particularly be at risk for Toxoplasma and cryptococcal infections, in addition to other infections seen at higher counts. While positive serology and rim-enhancing brain lesions that improve with appropriate therapy confirm the diagnosis of CNS toxoplasmosis, it is notable that toxoplasmosis more rarely causes cavitary pulmonary lesions, demonstrating that multiple opportunistic diseases can present at once. If improvements had not been seen, next steps would include brain biopsy for possible lymphoma or other infectious causes.
William Minteer Dr. R. Ramkissoon Dr. X.J. Wang	*Holy Hematopoiesis!* Case Presentation: A 70-year-old man with severe aortic stenosis recently diagnosed with systemic mastocytosis and chronic monomyelocytic leukemia presented for evaluation of a cholestatic liver injury. He was recently started on hydroxyurea for marked leukocytosis. His liver chemistries were previously within normal limits. On presentation, his alkaline phosphatase (ALP) vancomrtITJETQq163.58 69.024 356.35 265.7 reW*nBT/F1 11.04 Tf1 0

	Discussion: PTL is a rare malignancy accounting for less than 5% of all thyroid malignancies. Mantle cell lymphoma can present in the thyroid as an extra-nodal disease in only about 1% of cases. Furthermore, in cases of PTL presenting as an enlarging neck mass, only 1/3 of patients experience actual compressive symptoms of dysphagia, stridor and hoarseness. A high index of suspicion is needed to diagnose PTL, and core needle biopsies (rather than FNA) or excisional biopsy of an affected lymph node are needed to reliably diagnose lymphoma. Concern for PTL is raised in the setting of marked hypoechogenicity and rapid enlargement of the thyroid gland. Once diagnosed, chemotherapy is the treatment of choice for PTL.
Danny Mohama Dr. Antenneh Zewde	*Acquired Acrodermatitis Enteropathica and Other Signs and Symptoms of Nutritional Deficiency Following Bariatric Surgery* Introduction: Micronutrient deficiencies are common following bariatric surgery. These often include water-soluble vitamin, trace metal, and Vitamin D insufficiencies. Signs and symptoms of these deficiencies are numerous. However, a skin examination is an underutilized and important tool toward their identification. In the developed world, it is rare for these nutritional deficiencies to be present in severe and longstanding ways to cause dermatologic changes. Bariatric surgery patients remain at risk with restrictive and malabsorptive states, especially with difficulty adhering to proper nutritional supplementation. This case exhibits skin changes associated with multiple micronutrient deficiencies most notably acquired acrodermatitis enteropathica associated with zinc deficiency. Case Presentation: TM is a 45 year old woman with medical history of severe malnutrition, pancreatitis s/p total pancreatectomy and islet cell

	roles in human growth and development. The prevalence of zinc deficiency in the United States is between 1-3% and up to 20% worldwide. Deficiency is less common in developed countries due to its prevalence in dairy products, green leafy vegetables, and whole grains.
Jwan Naser Dr. Nandan Anavekar	*Left Ventricle Wall Rupture: Protected from Tamponade?* Case Presentation: A 86-year-old man presented with sharp central chest pain radiating up his neck associated with diaphoresis and dyspnea. He had a remote history of coronary artery bypass graft, atrial fibrillation and sinus node dysfunction requiring dual chamber pacemaker. On physical exam, there was no evidence of abnormal heart sounds or volume overload. ECG showed paced rhythm. Troponins were elevated and flat. He was admitted to the cardiac intensive care unit. Given improvement of chest pain and hemodynamic stability, an ischemia-guided approach was considered. Transthoracic echocardiography (TTE) showed ejection fraction of 45%, regional wall motion abnormalities mainly affecting the inferoseptal, anteroseptal and anterior walls, and a loculated pericardial fluid collection adjacent to the anterolateral left ventricular (LV) wall. Contrast TTE showed extension of the contrast agent from the main LV cavity to the adjacent pericardial fluid cavity, consistent with contained rupture of the LV wall or psuedoaneurysm. Cardiovascular surgery recommended computed tomography angiography of the thoracic aorta and coronary arteries which confirmed LV myocardial rupture along the anterolateral wall with active extrava complicated by acute kidney injury. In the setting of prior cardiac surgery and increased risk for dialysis, the patient and his family opted for a conservative approach in management. The

	systems, frequently presenting with hilar adenopathy, pulmonary infiltrates, and dermatologic manifestations. Less commonly, sarcoidosis can involve other organ systems, including ocular, cardiac, and hepatobiliary. Evaluation involves a comprehensive work-up to eliminate alternate diagnoses, while also gathering data to support sarcoidosis. Case Presentation: A 53-year-old male with rheumatoid arthritis on methotrexate and adalimumab, COPD, OSA, and non-alcoholic fatty liver disease, presented with a 3-month history of dyspnea, cough, acute arthralgias, and intermittent fevers. Prior to presentation, he sought care in the outpatient setting and was treated with multiple courses of antibiotics for his fever of unknown origin without improvement. Due to recurrent fevers to 104F and worsening dyspnea, he re-presented for evaluation, and was noted to have a new transaminitis, right axillary adenopathy, and asymptomatic hypercalcemia to 10.7 mg/dL. A Chest CT was performed, which revealed numerous enlarged axillary lymph nodes, diffuse parenchymal nodularity, fibrosis, and traction bronchiectasis, consistent with an infectious or inflammatory etiology. Infectious disease was consulted, and the patient underwent another broad infectious work-up, along with a bronchoscopy with bronchoalveolar lavage. This was unrevealing for a microbial source, gross airway abnormalities, or evidence of malignancy. Transthoracic echocardiogram showed no systolic or diastolic dysfunction. A work-up of hypercalcemia revealed an appropriately suppressed PTH, low 1,25-vitamin-D, normal ACE, and normal PTHrP. The hepatobiliary team was consulted to assist in the workup of his elevated alkaline phosphatase (peak 1025 U/L) and transaminitis (AST and ALT 100s). An MRCP revealed hepatosplenomegaly with mild fibrosis but was otherwise unremarkable. Axillary lymph node biopsy ultimately revealed granulomatous disease, suggestive of sarcoidosis. With the constellation of symptoms of adenopathy t with erythema nodosum, infectious workup and granulomatous findings on biopsy, he was diagnosed with sarcoidosis with hepatobiliary involvement. He was initiated on a steroid taper, with improvements in symptoms, normalization in liver enzymes and calcium, and resolution of fevers. Discussion: Sarcoidosis is a diagnosis of exclusion, as demonstrated by this case. It is imperative to initiate a broad infectious workup, rule-out other causes for granulomatous disease, and to screen patients for cardiac and ophthalmologic involvement due to risk of fatal or permanent impairments. Lofgren syndrome is a rheumatologic manifestation of sarcoidosis that presents with acute arthritis, bilateral hilar lymphadenopathy, and erythema nodosum. Presence of all three findings is 95% specific for Lofgren syndrome, and confirmatory biopsy is often not required for diagnosis of sarcoidosis. Though our patient did not have findings of erythema nodosum, his other presenting findings were consistent and not explained by an alternative cause after exhaustive workup.
Nathan Nelson Dr. Monazza Chaudhry	*A Case Where the Type of Testing Makes the Difference for Legionella* Introduction: Legionella pneumophila is a common cause of community

Dr. James Leatherman

acquired pneumonia (CAP). Legionella pneumophila can be difficult to diagnose because of multiple serogroups and subtypes not detected by the standard urine antigen test. This case describes the hospital course of a patient with CAP due to a legionella subtype with a negative urine antigen test and

Case Presentation: A 65-year-old female with a past medical history of multiple myeloma s/p chemotherapy and autologous stem cell transplant on

	Case Presentation: A 68-year-old man with a history of schizoaffective disorder was evaluated in the clinic for recurrent lightheadedness and pre-syncopal episodes with transient bradycardia for several months. EKG in the - from the clinic with a plan for outpatient workup. Given EKG abnormalities with reported symptoms, the patient was called back for close observation and further workup per cardiology recommendation. The patient was admitted with ongoing cardiac telemetry. Transthoracic echocardiogram did not reveal any significant structural abnormalities to explain clinical symptoms. On the second day, the telemetry revealed a 37 seconds episode of ventricular asystole due to a complete AV block, followed by a ventricular escape rhythm at approximately 30 beats per minute, later followed by the resumption of 1:1 AV conduction. The event was associated with a transient loss of consciousness, and an urgent temporary trans-venous pacemaker was placed. The patient subsequently received a dual-chamber permanent pacemaker. The patient was seen six weeks after discharge with no recurrence of lightheadedness or syncope. Discuss: -fascicular presyncope. Althou - without symptoms does not predict a high rate of progression to complete AV block, it is essential to note that when symptoms are associated with this EKG abnormality, there is an increased likelihood of progression to complete AV block, which could be intermittent. In one study by Marti-Almor et al., syncope was a predictor of progression to complete AV block in patients with tri-fascicular block. Management of patients presenting with syncope and tri-fascicular block involves clinical and electrophysiological evaluation of AV conduction system and consideration of a permanent pacemaker.
K e v i n O ô l Dr. Peter Lund	*Heartbreaker: A Case of AML Relapse Presenting as Myocardial Infarction* Introduction: Acute chest pain in young healthy individuals is unlikely to be due to cardiac ischemia. Nevertheless, clinicians must consider this pathology in the differential. Prompt diagnosis of ischemia can expedite treatment and avoid further sequalae. Case Presentation: A 21-year-old male with history of NPM1 mutated high

COVID-19 testing were negative. Hematology consulted and a bone marrow biopsy found a hypercellular bone marrow with 85% blasts. Treated with induction chemotherapy. EF remained 35-40% on subsequent imaging studies with development of left ventricular thrombus. After discharge, patient later underwent allogenic bone marrow transplantation from sibling with 6 month follow-up bone marrow showing no evidence of disease.

Conclusion: This case represents a rare presentation of acute myocardial infarction due to AML relapse in a young individual. A 2012 review found that only 12 of 774 (1.6%) cases of AML presented with myocardial infarction at time of diagnosis. Hyperleukocytosis causing leukostasis is proposed as one mechanism, however this was unlikely to be present in our case due to WBC on admission of 8.8. Other proposed etiologies in the literature include effects of prior chemotherapy, disseminated intravascular coagulation, hyperhomocysteinemia and deficiency of coagulation factors.
References:
1. Lukenbill, Joshua, et al. "Impact of Myocardial Infarction On Survival in Acute Myeloid Leukemia." Blood 120.21 (2012): 4321.
2. Chang, Hung, et al. "Acute myeloid leukemia associated with acute myocardial infarction and dural sinus thrombosis: the possible role of leukemia-related hyperhomocysteinemia." Journal of the Chinese Me1(em)5(i)-

	biopsy results, he was started on vancomycin and Zosyn empirically. Anti-fungal treatment was deferred. On ROS, he denied any dysphagia, odynophagia or shortness of breath. He reported abdominal pain associated with eating, early satiety and a 20 lbs weight loss over the preceding 4 months. He reported recent travel throughout US, Jamaica, Costa Rica, Puerto Rico, and Italy. He denied any other relevant exposures or risk factors. Initial laboratory survey was notable for HGB 7.8 (N 13.1-17.5), elevated alkaline phosphatase 234 (N 40-129) with BMP, CBC and LFTs otherwise unremarkable. Bacterial and fungal blood cultures, Cryptococcal antigen, urine histoplasma and sputum AFB were all negative. CMV IGG was positive. Bronchoscopy was performed and returned negative for pneumocystis with no growth otherwise. An inguinal lymph node biopsy was performed without immunophenotypic evidence of non-Hodgkin lymphoma. Imaging studies including CT neck revealed subcutaneous soft tissue nodules throughout the head, face, and neck with prominent cervical lymph nodes and tonsillar hypertrophy concerning for disseminated fungal infection or IRIS. CT chest, abdomen and pelvis also noted diffuse skin and subcutaneous nodules particularly in the abdomen and pelvis as well as multiple enhancing nodules in the lungs, pleural space and thoracic paraspinal area with extensive mesenteric and retroperitoneal adenopathy. Skin biopsy returned positive for HHV8 and CD34 on immunohistology stains confirming a diagnosis of Kaposi sarcoma. CD68 was negative ruling out a granulomatous inflammation. Acid fast, GMS and PAS special stains were negative ruling out disseminated mycobacterial and fungal infection respectively. Conclusion: Commonly implicated causes of IRIS including mycobacterial, cryptococcus, pneumocystis, cytomegalovirus, herpes, hepatitis B and C were ruled out further supporting the diagnosis of IRIS-like syndrome in our patient. It is postulated that IRIS occurs in >10% of patients with KS, tuberculosis, or cryptococcus infection and visceral involvement leads to considerable morbidity and mortality (Achenbach). We present this case to increase awareness of IRIS-like syndrome and stress the importance of maintaining close clinical supervision in patients with KS while initiating, changing or resuming HAART (Leidner).
Jessica Park	*Small Bowel Obstruction Caused by Carcinoid Tumor* Conclusion: As cancer therapies advance to personalize treatment and target specific immune responses, it is important for providers to be aware of the potential adverse effects such as the potential for increased production of PTHrP leading to hypercalcemia.
Nisha Patel Dr. Angela Grassi Dr. Mariana Restrepo-Holguin Dr. Sunnia Chen	

	Case Description: A 47-year-old male sanitation worker presented with a two-week history of fevers, right upper quadrant pain, and jaundice. Labs were significant for elevated liver enzymes in a hepatocellular pattern and thrombocytopenia. Imaging revealed sludge in the gallbladder with mild wall thickening, diffuse hepatic steatosis, and multiple wedge-shaped hypoenhancing areas in the spleen suggestive of infarcts. The patient was admitted due to concern for acute cholangitis and incidentally found splenic infarcts. During hospitalization, MRCP showed no biliary ductal dilation or filling defects, which ruled out acute cholangitis, and TEE was negative for evidence of infectious endocarditis. The patient continued to have elevated liver enzymes and persistent fevers despite antibiotic coverage with piperacillin-tazobactam. Additional workup was significant for elevated antiphospholipid antibodies in the setting of splenic infarcts. With no clear broadened to include a variety of atypical infectious causes of acute hepatitis. Liver biopsy subsequently revealed non-necrotizing lobular and fibrin granulomas. Due to suspicion for Q fever infection, the patient was empirically started on doxycycline with subsequent clinical improvement. Coxiella burnetii titers and PCR resulted positive in a manner consistent with acute Q fever infection, and the patient was started on hydroxychloroquine as well. His hospitalization was also complicated by upper GI hemorrhage secondary to anticoagulation in the setting of positive antiphospholipid antibodies. EGD was performed, revealing four duodenal ulcers and an actively bleeding pre-pyloric ulcer, which was clipped. Discussion: This case highlights the importance of considering a broad differential diagnosis, including Q fever, in patients with signs of hepatobiliary disease. Q fever has a variable presentation and can often resemble other infections or autoimmune diseases. While hepatitis is commonly associated with Q fever, acalculous cholecystitis is a very rare presentation. In addition, although transient antiphospholipid antibodies are a known phenomenon of Q fever, they are rarely associated with independent thrombotic complications in patients without prior diagnosis of antiphospholipid antibody syndrome or evidence of underlying Q fever endocarditis. However, our patient suffered an extensive thrombotic event necessitating anticoagulation, despite relatively low level positivity of anticardiolipin IgG (60.9). Our findings suggest that patients with Q fever should be routinely tested for antiphospholipid antibodies, with further workup for thrombosis and consideration of anticoagulation as needed. Furthermore, due to the significant morbidity Q fever patients may suffer secondary to thrombosis, close follow-up with vascular medicine specialists is warranted.
Marissa Paulson Dr. Abdulsabur Sanni	*Initial Concern for One in a Million Vaccine Side Effect leads to Diagnosis of Acute Promyelocytic Anemia* Introduction: When this 41 year old patient began developing bruising around her abdomen and her typical migraines increased in severity, her initial concerns were not for her eventual diagnosis of cancer, but of an exceedingly rare diagnosis that was being splashed across media at the time- thrombotic thrombocytopenia. This patient had received the Johnson and Johnson (J & J) Ad26.COV2.S vaccine against severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) two months prior to symptom onset. She presented to care

recipients but was eventually diagnosed with Acute Promyelocytic Leukemia (APML).

Case presentation: This 41-year-old woman presented to clinic complaining of fatigue, headaches, dyspnea with exertion, vaginal bleeding, and bruising over her abdomen and bilateral lower extremities. She had received the J & J

Hemoglobin on post-hospital follow up 5 days later was 5.7 g/dL, prompting re-admission to the outside hospital. Despite repeated blood transfusions (total of 20 units), the anemia persisted. Extensive imaging to look for indolent bleed including endoscopy, colonoscopy, tagged red blood cell study, CT angiography of abdomen/pelvis/lower extremities, CT chest/abdomen/pelvis, MRI right femur, and repeat lower extremity ultrasound with doppler was unrevealing. Labs including infectious studies, hemolysis labs, peripheral smear, and bone marrow biopsy were also inconclusive. He was subsequently transferred to our institution for further workup/management of refractory anemia.

Upon transfer, Rheumatology was consulted for comprehensive evaluation. During our initial exam, corkscrew hairs, perifollicular hemorrhages, and purpura were appreciated, raising suspicion for scurvy and prompting us to obtain an in-depth dietary history. He described a diet completely devoid of fruits or vegetables and reported eating only one meal per day prior to hospitalization, which was frozen and processed. Otherwise, he reported snacking almost exclusively on chips. A vitamin C level was checked and returned undetectably low. Dermatology and Hematology/Oncology were also consulted and agreed that scurvy was the most likely diagnosis. He was treated with intravenous and oral ascorbic acid with significant improvement in cutaneous findings and no further need for blood transfusions.
Conclusion: This case highlights the importance of a thorough physical exam and history in reaching a diagnosis. Albeit rare, scurvy does still occur in

abdomen and pelvis showed pan colitis after which patient was started on oral vancomycin and metronidazole for suspected CDI. Patient did develop diarrhea following admission and a stool sample returned positive for Clostridium Diffcile. General surgery was consulted and discussed surgical intervention which included total colectomy with end ileostomy for fulminant CDI. After this discussion, it was felt that the patient would not want to

	that typically affects young women in their 30s to 40s. Patients most often present with signs and symptoms of cystic lung disease including pleural effusions, spontaneous pneumothorax, and progressive dyspnea with exertion. Other symptoms can be associated with extrapulmonary sites related to renal angiomyolipoma or lymphatic vasculature including chylothorax. Diagnosis of LAM is typically a clinical diagnosis which includes characteristic thin-walled, well defined, diffusely distributed pulmonary cysts, VEGF D level>800 pg/ml, chylothorax, and renal angiomyolipoma. Definitive diagnosis is made with lung biopsy via VATS. There is currently no cure for LAM, but mTOR inhibitors like Sirolimus can help slow the progression of the disease.
Sumar Quint Dr. Samuel Ives	*Hold Your P: Renal Phosphorous Wasting, a Rare Side Effect of IV Iron Infusion* Introduction: Iron deficiency anemia (IDA) is a prevalent disease, affecting up to 20% of women worldwide. Treatment with parenteral iron is common due to oral iron side effects and absorption difficulties. An underrecognized but serious side effect of a common IV iron formulation, ferric carboxymaltose (FCM), is hypophosphatemia. This adverse effect is thought to be caused by increased circulating levels of fibroblast growth-factor-23 (FGF-23) which leads to inappropriate renal excretion of phosphorous. Risk factors for developing severe hypophosphatemia after FCM therapy include patients who are chronically malnourished or have malabsorption syndromes. Case Presentation: A 36-year-old female with past medical history of bulimia nervosa and IDA presented to the emergency department for chest pain, she was found to have phosphorous of 1.0 initially thought to be due to malnourishment and refeeding syndrome. She was briefly admitted and given oral replacement. The patient re-presented two days later after labs at a residential eating disorder facility showed continued hypophosphatemia. Symptoms included headache and muscle weakness. In the ED th phosphorous was 1.4, she was admitted for IV repletion and further work up. Parathyroid hormone (PTH), 25-OH Vitamin D level, and ionized calcium were within normal limits. The patient did not have any other persistent electrolyte disturbances. phosphorous was continually repleted, with only transient improvements in serum levels. A 24-hour urine phosphorous was found to be high at 1718 mg/day (400-1300 mg/day), suggesting renal wasting of phosphorous. Given no other endocrine or lab abnormalities, prior medications were investigated. The patient had been seen by hematology three weeks prior to admission and given two doses of FCM infusion for IDA. The suspicion of FCM causing phosphorous wasting was confirmed with an inappropriately normal FGF-23 level of 122. The patient was discharged with a PICC line for outpatient infusion of phosphorous, and follow-up with endocrinology. Discussion: FCM infusions are thought to cause hypophosphatemia in susceptible patients by increasing FGF-23 levels by a mechanism that is not fully understood. FGF-23 is a bone-derived hormone that inhibits phosphate transporters in the renal tubules and decreases activation of 1,25-OH Vitamin D, leading to both renal wasting and inability to absorb phosphorous. Risk factors for hypophosphatemia secondary to FCM include malnourishment, hyperparathyroidism, malabsorption syndromes, and vitamin D deficiency. In the case presented the patient had a long history of disordered eating and was likely deficient in total body phosphorous before the FCM infusion. The side effect of severe hypophosphatemia should be better known to clinicians, as

	patients with risk factors for hypophosphatemia are often the most common patients to have IDA. Treatment consists of frequent lab checks and replacement of phosphorous and most cases resolve within 3 months.
Alison Raffman	*Non-Cardiogenic Pulmonary Edema Following Naloxone Use for Heroin Overdose* Introduction: Naloxone is an opioid antagonist medication that is commonly used as a reversal agent in cases of opioid overdose. This medication is generally considered as safe to give for this purpose, however it has been associated with several uncommon complications. Non-cardiogenic pulmonary edema following Naloxone administration has been reported as a rare adverse event associated with use of this medication. Case Presentation: A 32-year-old male with prior history of opioid abuse presented to the emergency department after reported IV heroin overdose the patient unresponsive at home, called EMS and initiated bystander CPR at the scene. She also administered 1.2mg IM Naloxone with subsequent immediate improvement in patient subsequently developed worsening shortness of breath, was found by medics with oxygen saturations in the 80s, and required oxygen via non-rebreather en route to the hospital. On arrival to the emergency department the patient was alert and oriented, reporting shortness of breath but otherwise without complaints. Significant initial vitals included HR 129, RR 29, oxygen saturation 89%, the patient was otherwise normotensive and afebrile. His exam was notable for coarse breath sounds bilaterally (greater on the left). Bedside cardiac ultrasound did not demonstrate any significant abnormalities, and lung ultrasound demonstrated B-line predominance bilaterally. Chest x-ray demonstrated bilateral infiltrates (moreso on the left). CT chest again showed bilateral infiltrates that were more prominent on the left, consistent with asymmetric pulmonary edema. The patient was monitored in the emergency department and was eventually transitioned from non-rebreather to 3L O2 via nasal cannula, however after four hours of monitoring the patient was trialed on room air and experienced worsening tachypnea with desaturation to 90%. The patient was admitted to medicine for observation, and after several hours the patient was successfully transitioned to room air without significant hypoxia or increased work of breathing, with clear breath sounds noted on subsequent examinations. The patient was able to be discharged home in good condition. Discussion: Non-cardiogenic pulmonary edema has been previously reported as a rare complication of Naloxone administration following opioid overdose. pathophysiology of pulmonary edema in such cases, however a dose-response relationship between Naloxone and subsequent pulmonary complications has been described previously. Overall, this case highlights a case of respiratory distress associated with non-cardiogenic pulmonary edema following Naloxone administration for opioid overdose, and demonstrates successful management of this condition(i)-4(s)9(t)-4(o04 Tf1 0 0 G[(r)-3(ay d)9(em)-5(o)

device. Complications of endocarditis are vast and include cardiac valve destruction, perivalvular extension, systemic embolization, metastatic spread of infection, and immunologic manifestations[1]. Infective endocarditis also carries a high mortality rate. In hospital mortality rate for patients with infective endocarditis is 18-23%, and the six-month mortality rate of patients with infective endocarditis is 22-27% [2-4]. Given the significant complications and high rate of mortality of infective endocarditis, prevention and prompt treatment carry the upmost importance.

Case Description: The patient is a 61-year-old male who presented with 10 days of weakness, myalgias, increasing dyspnea on exertion, and fevers. Relevant medical comorbidities include severe aortic stenosis secondary to bicuspid aortic valve status post aortic valve replacement with a bioprosthetic valve eight years prior. On physical exam, a III/VI systolic ejection murmur was appreciated at the right upper sternal border. There were no peripheral manifestations of endocarditis or embolic phenomenon appreciated on physical exam. A transesophageal echocardiogram showed a 3 mm echodensity on the anterior leaflet of prosthetic valve and thickening of the mitral-aortic intervalvular fibrosa. A CT cardiac angiogram showed a para-aortic phlegmon. Blood cultures grew Granulicatella adiacens, a streptococci species. He initiated intravenous vancomycin and ceftriaxone for presumed endocarditis. The patient then underwent surgical debridement of previously implanted aortic valve and entire root complex and aortic root replacement utilizing a 25-mm On-X composite valved conduit. Extensive purulent material was noted intraoperatively. He continued vancomycin for six weeks then transitioned to oral penicillin for suppression for at least 12 months.

Discussion:
put him at risk of infective endocarditis. The significant mortality of infective endocarditis makes this diagnosis a medical emergency requiring prompt hospitalization, acquisition of blood cultures, empiric antibiotic therapy, and expedited cardiac and systemic imaging to confirm the diagnosis and evaluate for complications. Granulicatella is a nutritionally variant streptococci, meaning it grows best on media supplemented with thiol or pyridoxal [5]. It is a catalase-negative, oxidase-negative, facultative anaerobe and is normally present in oral flora and rarely manifests as invasive infections [6]. Interestingly, Granulicatella has only been implemented in up to five percent of cases of streptococcal endocarditis [5]. Targeted antimicrobial therapy is essential to ensure elimination of bacteremia and treatment of infective endocarditis.
References:
1.Mansur,A.J.,etal.,TheComplicationsofInfectiveEndocarditis:AReappraisalin the1980s.ArchivesofInternalMedicine,1992.152(12):p.2428-2432.
2. Hasbun,R.,etal.,ComplicatedLeft-SidedNativeValveEndocarditisinAdultsRiskClassificationforMortality.JAMA, 2003.289(15):p.1933-1940.
3. Hill,E.E.,etal.,InfectiveEndocarditis:changingEpidemiologyandPedictors of 6-
2006. 28(2): p. 196-203.
4. Wang, A., et al., Contemporary Clinical Profile and Outcome of Prosthetic Valve Endocarditis. JAMA, 2007. 297(12): p. 1354-1361.
5. Gupta, S., et al., Granulicatella adiacens abscess: Two rare cases and

Case Presentation: A 78-year-old male presented to the Emergency Department with two-days of fevers, nausea, vomiting, confusion, and

nephrostomy tube was discussed but ultimately deferred due to an atrophic right kidney with presumed trivial function and a decompressed ureter. The patient was discharged home with outpatient oncology follow up scheduled in one month. During her oncology appointment, she described increased right thigh pain and swelling, otherwise denied new symptoms. Staging PET scan showed rupture of the upper pole of the right kidney due to hydronephrosis from progressive malignant obstruction complicated by an extensive urinoma extending from the posterior right kidney along the retroperitoneum and psoas/iliopsoas musculature into the muscles of the proximal medial right thigh. She was readmitted to the hospital, and these findings were confirmed with a CT Urogram with the urinoma measured to be 36 cm. Percutaneous drains were placed to the right perinephric, right retroperitoneal, and right upper leg fluid collections. Nephrostomy tube was deferred due to urinoma infection with E. coli. A CT urogram was repeated 48 hours after tube placement and showed interval decrease in all three fluid collections. She was discharged home on antibiotics with plans for future nephrostomy tube placement. She was initiated on radiation therapy to shrink the tumor and relieve obstruction.

Discussion: This is a rare case of locally advanced cervical cancer causing ureteral obstruction leading to hydronephrosis, calyceal rupture, and an extensive urinoma presenting as right thigh pain. RCR is more commonly seen with acute renal stone impaction, and is less common in malignancy, potentially secondary to the subacute/chronic obstruction from tumor burden. Treatment of urinoma includes percutaneous drain placement with or without nephrostomy tube to decompress the kidneys. Nephrostomy tube placement

Discussion: While Amphotericin B is the main therapy to treat blastomycosis in severe infections. The robust inflammatory response that accompanies severe infection has led clinicians to utilizing corticosteroids. For patients with severe respiratory failure such that they cannot adequately ventilate or oxygenate their lungs while on mechanical ventilation, ECMO has been utilized. Corticosteroids and ECMO are logical choices in severe disease as these therapies are utilized in other settings of ARDS. It remains unclear as to the efficacy of these therapies in reducing morbidity or mortality as randomized control trials have not been conducted.

	unifying clinical diagnosis of IgAV and initiating early surveillance for sequelae and complications.
Mackenzie Stice	: K H Q : H D N Q H V V L V Q ¶ W ' H F R Q G L W L R Q L Q J

Wintheiser
Dr. Mohammed Bhuiyan

Introduction: Immune-mediated necrotizing myopathy (IMNM) is a rare inflammatory myopathy distinguished on pathology by necrotic muscle fibers without significant inflammation. It is associated with anti-3-Hydroxy-3-Methylglutaryl-CoA Reductase (HMGCR) and anti-signal related peptide (SRP) antibodies, although up to 20% of patients are seronegative. Like other immune-mediated myopathies, it presents with progressive muscle weakness and elevated serum creatine kinase (CK). Treatment consists of immunosuppression with some combination of steroids, immunomodulators, intravenous immunoglobulin (IVIG), methotrexate, and rituximab.

Case Presentation: A 78-year-old woman presented with four days of bilateral lower extremity weakness. Physical exam revealed proximal muscle weakness in the deltoids and hip flexors. Labs demonstrated a CK of 26,000 unit/L and marked transaminitis (ALT 617, AST 1338). She also had a significant AKI (creatinine 10.51) and ultimately required dialysis. Notably, she had been started on statin therapy several months prior. This, combined with her

	relative hypoxic environment further augmenting production of lactic acid. Decreased lactate utilization from extensive liver metastasis might also play a role in some cases. Malignancy-associated lactic acidosis is associated with a very high mortality rate and is usually fatal without the initiation of treatment for underlying malignancy. We report 6 selected cases of tumor-associated lactic acidosis, and describe 3 in detail that demonstrate interesting pathophysiology related to the condition. Localization of sites of production and excretion of lactic acid are presented in the case of a 71-year-old man with metastatic small cell lung cancer. Chronic hypokalemia and cancer cachexia are observed and discussed in our second and third cases of metastatic leiomyosarcoma. Lactic acid metabolism and the treatment of lactic acidosis in neoplastic disease are also discussed. The observation of this number of cases of lactic acidosis in solid malignancies in one practice suggests the syndrome is more common than appreciated. Awareness of the syndrome is critical to provide appropriate care.
Benjamin Swart Dr. Brooke McDonald	+LFNDP¶V 3XUSXUD" 0XOWLSOH 9DVFXODLW Introduction: The presence or absence of immune complexes is one of the major differentiating factors of acute glomerulonephritis (GN). Lupus nephritis is one of multiple GNs characterized by immune complex deposition. Pauci-immune vasculitides, in contrast, are characterized by a lack of immune complex deposition. This case report describes a patient with rapidly progressive GN secondary to both types of mechanisms. Case Presentation: A 67-year-old male with past medical history pertinent for lupus nephritis, CKD 3, hypertension, and hypothyroidism presented with five days of altered mental status and fever. Vitals at presentation were within normal limits. On exam he was obtunded, with a non-blanchable, petechial rash on his lower extremities, bilateral myoclonus in upper extremities, and cola-colored urine. Initial labs were notable for WBC count 24.51 cells/mm2, hemoglobin 11 g/dL, and platelets 302 cells/mm2. Basic metabolic panel revealed serum potassium 5.4 mEq/L, BUN 33 mg/dL and creatinine of 3.94 mg/dL (baseline of 1.5). Inflammatory markers were elevated with CRP of 225.7 mg/L and ESR of 46 mm/hr. Complements 3 and 4 were within normal limits. HIV, hepatitis B/C and cryoglobulins were negative. IgA and IgG were within normal limits and CSF was unremarkable. His urinalysis was notable for proteinuria and hematuria with no growth on urine culture. Head CT and renal ultrasound were both within normal limits. Lupus anticoagulant was positive. Anti-neutrophil cytoplasmic antibody (ANCA) was negative, however myeloperoxidase (MPO) antibody was strongly positive (46 AU/mL). With low suspicion for active infection, the patient was started on high dose corticosteroids and emergent dialysis in the setting of worsening uremia. Kidney biopsy was performed and demonstrated focal proliferative lupus glomerulonephritis (classes III and V) in addition to necrotizing small artery vasculitis consistent with MPO/ANCA vasculitis. Skin biopsy was positive for leukocytoclastic vasculitis without lupus involvement. The patient was started on monthly cyclophosphamide with mesna infusion for dual treatment of both lupus nephritis and ANCA vasculitis and hydroxychloroquine with prednisone taper. By the time of discharge, his kidney function had significantly improved.

	Discussion: Though rare, there are case reports of an overlapping syndrome of systemic lupus erythematosus and ANCA/MPO-associated vasculitis. It usually presents in women in the fourth decade of life and diagnosed via kidney biopsy. Treatment is not standardized, but can include cyclophosphamide, rituximab, hydroxychloroquine, plasma exchange, IV immunoglobulin, and steroids. When presented with conflicting lab values, be open to the fact that a single unifying diagnosis may not exist. In the search can have as many di
Reema Tawfiq Dr. Christopher Tipton Dr. Akeem Lewis Dr. Naseema Gangat **2021 MN-ACP Resident Peopleôs Choice Award**	7 U L S O H 1 H J D W L Y H 0 \ H O R I L E U R V L V Z L W K 3 *Syndrome* Introduction: Myelofibrosis is a myeloproliferative neoplasm characterized by marrow fibrosis and extramedullary hematopoiesis. The median age of presentation is 67 years. In rare occurrences, it can affect younger adults. Commonly implicated driver mutations include JAK2, CALR and MPL. Triple-negative MF makes up 10-15% of patients with MF and lends to a higher rate of leukemic transformation and poor nBT/n(c)9(t)-4(r)7(an)9(sf)-5(

	adrenal insufficiency and hemosiderosis.

Rebecca Yao